



BRENT D. CARLSON, MD
Orthopedic Surgeon
www.carlsonmd.com

1200 N 10TH ST. W. STE A
ALTOONA WI 54720
TEL 715.832.1400

757 LAKELAND DR. STE B
CHIPPEWA FALLS WI 54729
TEL 715.723.8514

Post-op Instructions for Shoulder Capsular Repair, Bankardt, or SLAP Repair

These instructions are to compliment the information given by the nursing staff and physical therapists. They cover many of the common questions.

Wound Care

- Dressings are to be kept clean and dry. You may change the dressing daily if desired. After removing the old dressing replace with 4x4 gauze pads and secure with tape. A small amount of clear drainage or bleeding is normal. If this is happening, the dressing should be changed daily.
- You may get the incision wet when showering 3 days after surgery. The shower should be brief and the wound patted dry with a clean towel. No baths or soaking the incision until 2 weeks after surgery and scabs are absent. It may be comfortable to use a rolled up towel as a pillow under your arm while showering. You may also bend over at the waist and gently dangle your arm to wash underneath it.
- If purulent drainage (thick white or greenish in color) is coming from the wound, or the wound has increasing redness, or if you are having a temperature of 101, please report these symptoms to your surgeon or the doctor on call.
- Never, ever remove your own stitches or trim what may appear to be excess suture material. We will remove your stitches in the office at your post-op appointment. If you are concerned about your stitches or if they are bothering you, please call us.

Pain

- Ice your shoulder as frequently as possible. We recommend 4 - 5 times per day for 20 minutes per time. You may use either the ice bag/cooling apparatus given to you at the hospital or simply place ice in a zip lock bag and place on the shoulder.
- Narcotic pain medication will be prescribed for use after you leave the hospital. Try to wean down as tolerated. These medications can cause constipation and you may want to use an over the counter stool softener. Tylenol products may be used instead of the prescribed pain medication. The prescribed narcotic medications should not be taken at the same time as plain Tylenol. You may supplement your pain medication by using ibuprofen or Aleve for any breakthrough pain. It can help to stagger your pain medication with ibuprofen or Aleve as needed. If a refill of medication is needed, please call the office during regular business hours, Monday-Friday 8:00 a.m. to 5:00 p.m. In general, refills will not be made after hours or on weekends, so please plan ahead.

Driving

- To drive you must no longer be taking narcotic pain pills (plain Tylenol is allowed). Also, you must feel strong and alert. You may take your arm out of the sling to hold the bottom of the steering wheel; you should NOT actively raise your arm until cleared by physical therapy. Most people start driving approximately two weeks after surgery.

Exercises

- Leave your arm in the sling until you are told otherwise by your surgeon or physical therapist. You may take it out of the sling for range of motion exercises and showering.
- Upon discharge from the hospital you are encouraged to perform hand, wrist and elbow range of motion exercises 4 - 5 times per day. These exercises will help to decrease swelling. You should not attempt to elevate the surgical arm under its own muscle power. Your physical therapist will progress your activity appropriately.

Follow-up

Make sure an appointment has been scheduled for you at Chippewa Valley Orthopedics & Sports Medicine for approximately 1 - 2 weeks after surgery.

Brent Carlson, M.D.
Chippewa Valley Orthopedics & Sports Medicine
(715) 832-1400